

The role of angiogenesis in prostate and other urologic cancers: a review

Review

Synthèse

From the Departments of Urology and Cancer Biology, The University of Texas M. D. Anderson Cancer Center, Houston, Tex.

This article has been peer reviewed.

CMAJ 2001;164(5):662-70

Jonathan I. Izawa, Colin P.N. Dinney

Abstract

ANGIOGENESIS IS A PROCESS CRITICAL TO both tumour growth and metastasis. It is a dynamic integrated process involving basement membrane degradation, endothelial cell proliferation and migration, and capillary tubule formation. Under normal circumstances, the microvasculature is maintained in a quiescent state. The acquisition of the angiogenic phenotype depends on the outcome of stimulatory and inhibitory regulation by the tumour and its microenvironment. There are markers of angiogenesis that potentially could provide prognostic information in addition to that gained from conventional clinicopathologic data, and antiangiogenic therapy for urologic cancers has potential advantages over current therapeutic strategies. Promising preclinical studies have led to the initiation of phase I studies of antiangiogenic therapy in combination with chemotherapy, which may lead to novel treatments for urologic malignant tumours and may identify new intermediate markers for the response to therapy.

Prostate adenocarcinoma (PCa) is estimated to be the most commonly diagnosed noncutaneous malignant disease and the second leading cause of cancer deaths in men, whereas bladder transitional cell carcinoma (TCC) and renal cell carcinoma (RCC) are estimated to be the sixth and ninth most commonly diagnosed malignant diseases respectively.¹ Despite significant improvements in local and systemic therapies, most deaths from cancers are due to metastases that resist conventional therapies.²⁻⁴ A major barrier to effective treatment is the biologic heterogeneity of cancer cells exhibited by genetic, biochemical, immunologic and biologic characteristics, such as cell-surface receptors, enzymes, karyotypes, cell morphology, growth parameters and drug resistance. Because of this heterogeneity, continued empiricism in the treatment of urologic cancers is unlikely to produce significant clinical improvements in patient outcomes.³ Rather, treatments directed toward critical processes that are involved in tumorigenesis and metastasis are more likely to result in significant advances in the treatment of urologic cancers. There are several lines of evidence that carcinogenesis is a multistep process reflecting genetic alterations that occur in the transformation of normal cells into highly malignant derivatives.⁵ Similarly, the process of angiogenesis appears to be acquired in discrete steps during tumour development and is critical to tumorigenicity and metastasis.⁵ In this article, we review the compelling evidence that supports the importance of angiogenesis and its clinical relevance in urologic cancers.

Angiogenesis

Tumour growth and metastasis depend upon the development of a neovasculature in and around the tumour.⁶⁻¹⁰ This process, called angiogenesis, is regulated by the balance between stimulatory and inhibitory factors released by the tumour and its microenvironment (Table 1).^{8,11-15} Angiogenesis facilitates progressive tumour growth by providing adequate oxygenation to the tumour through a series of inter-related steps, including endothelial cell proliferation, motility of endothelial cells through the extracellular matrix toward angiogenic stimuli, and capillary differentiation (Fig. 1).

Much of our knowledge of the biology of PCa, TCC and RCC arises from animal experiments in which human tumour cells have been implanted into immune-

deficient nude mice. The angiogenic, tumorigenic and metastatic potential of these human xenografts grown in nude mice is enhanced by implanting these tumour cells and growing them within the orthotopic site rather than a heterotopic site.¹⁶⁻²⁰ Thus, mechanisms regulating angiogenesis are tissue specific,²¹ and the angiogenic phenotype is regulated by the differential expression of cytokines and growth factors within the microenvironment of the organ.²² This concept of organ-specific angiogenesis is important for the interpretation of preclinical studies evaluating angiogenesis and antiangiogenic therapies for human cancers, because heterotopic tumour models will not reflect accurately the interactions between the microenvironment of the organ and the tumour.

Prognostic markers

A biomarker is biologic material that can be used to enhance the detection of disease or provide prognostic information. The ideal prognostic biomarker would be noninvasive and would provide predictive information regarding the natural history of the disease or the response to treatment in addition to that gained from conventional clinicopathologic parameters. Markers of angiogenesis, including angiogenesis factor expression and microvessel density (MVD), have been evaluated as prognostic markers for urologic cancers.²³

Antiangiogenic therapies

Because tumour growth and metastasis depend on angiogenesis, a great deal of attention has been focused on therapy that can interrupt this process. Antiangiogenic therapy can target endothelial cells directly; inhibit the production or action, or both, of proangiogenic peptides by the tumour cells or host; or enhance the expression of angiogenesis inhibitors within the tumour.²⁴ Antiangiogenic therapy that targets endothelial cells, rather than tumour cells directly, has been evaluated as a novel therapeutic strategy for malignant diseases.^{25,26} The theoretical advantage to this therapy is that endothelial cells are unlikely to acquire mutations that lead to drug resistance.²⁷ Endothelial cells are also readily exposed to bloodborne agents, circumventing the problem of drug delivery. Although it was widely assumed that antiangiogenic therapy was antiproliferative, it has recently been found that antiangiogenic therapy can induce apoptosis and tumour regression.²⁸

Prostate cancer

Angiogenesis

Early studies reported that conditioned medium from cultures of human PCa cells stimulated endothelial cells, suggesting that these cells produce proangiogenic factors,²⁹ and both basic fibroblast growth factor (bFGF) and vascular endothelial growth factor (VEGF) proteins have been measured in the urine of patients with PCa.³⁰ Subsequently, it was reported that human PCa produces VEGF and interleukin-8 (IL-8) protein.³¹ These and other proangiogenic factors are believed to enhance the tumorigenicity and metastasis of PCa, because xenografts of the highly metastatic PC-3M-LN4 PCa cell line overexpressed bFGF, IL-8 and matrix metalloproteinase-9 (MMP-9)

Table 1: Factors implicated as angiogenic promoters and inhibitors in prostate, bladder and kidney cancers

Cancer	Promoter	Inhibitor
Prostate	Basic fibroblast growth factor (bFGF)	Endogenous inhibitor Interleukin-10 (IL-10)
	Vascular endothelial growth factor (VEGF)	Tissue inhibitor of metalloproteinase-1
	Interleukin-8 (IL-8)	Angiostatin
	Matrix metalloproteinase-9 (MMP-9)	Prostate-specific antigen
	Matrix metalloproteinase-2 (MMP-2)	Interferon
	Transforming growth factor- β_1	Pharmaceutical agent Anti-VEGF antibody (A4.6.1)
	Platelet-derived endothelial growth factor	Fumagillin analogue, 0-(chloroacetyl-carbamoyl) fumagillol (TNP-470)
	Nitric oxide	Linomide
		Carboxyamido-triazole
		Diet Energy-reduced diet Soy products
Bladder	bFGF	Endogenous inhibitor Thrombospondin
	VEGF	Interferon
	Midkine	Pharmaceutical agent Antiepidermal growth factor receptor antibody (C225)
	Thymidine phosphorylase	DC101
	Hepatocyte growth factor	Halofuginone
	Hyaluronic acid fragments	Diet Soy products
	IL-8	
	Transforming growth factor- β_1	
	Hyaluronidase	
	MMP-9	
Angiogenin		
Kidney	bFGF	Endogenous inhibitor Endostatin
	VEGF	Interferon
	Thymidine phosphorylase	Vitamin D ₃
	MMPs	Pharmaceutical agent Fumagillin analogue, 0-(chloroacetyl-carbamoyl) fumagillol (TNP-470)
	Urokinase-type plasminogen activator (uPA)	
	Transforming growth factor- β_1	
	Hepatocyte growth factor	

mRNA compared with the poorly metastatic parental PC-3M cell line from which it was selected.³² Similarly, the highly metastatic LNCaP-LN3 cell line overexpressed VEGF compared with either the parental LNCaP cell line or LNCaP-Pro5, which was selected for its tumorigenicity in the prostate.³³ LNCaP-LN3 was also more metastatic and overexpressed VEGF mRNA and protein and VEGF receptor protein, with enhanced tumour-induced angiogenesis, compared with either LNCaP or LNCaP-Pro5, after implantation into the prostate of athymic nude mice.

The importance of IL-8 as a mediator of PCa angiogenesis was shown by experiments in which PC-3P cells with lower metastatic potential were transfected with sense IL-8, whereas highly metastatic PC-3M-LN4 cells

were transfected with antisense IL-8 constructs.³⁴ Following orthotopic implantation, the sense-transfected PC-3P cells overexpressed IL-8 and MMP-9 and were highly angiogenic, tumorigenic and metastatic compared with parental PC-3P cells or controls. Antisense transfection of the PC-3M-LN4 cells reduced the expression of IL-8 and MMP-9 and tumour-induced angiogenesis, resulting in inhibited tumorigenicity and metastasis. Therefore, IL-8 also seems to be involved in the regulation of angiogenesis and tumour growth in PCa, in part by induction of MMP expression.

The angiogenic and metastatic phenotype of PCa was also shown to be directly related to the expression of transforming growth factor- β_1 and MMP-2 and inversely related




Fig. 1: Angiogenesis in (A) superficial and (B) muscle-invasive transitional cell carcinoma of the bladder is, in general, illustrative of the process of tumour neovascularization in other cancers. The multiple, sequential and interrelated steps by which angiogenesis is hypothesized to occur involve angiogenic stimuli or lack of endogenous angiogenic inhibitors, or both; the inciting by endothelial cells within venules of local proteolysis to degrade the basement membrane; the protrusion of endothelial cells through the wall of the venule; degradation of the interstitial matrix; continuing movement of endothelial cells toward the angiogenic stimulus; formation by endothelial cells of capillary sprouts that form a lumen; proliferation within sprouts; joining of tips of sprouts; blood flow; and formation of new basement membrane and incorporation of microvascular pericytes. TCC = transitional cell carcinoma, BM = basement membrane.

to the expression of IL-10, which upregulated a natural tissue inhibitor of metalloproteinase-1.^{35,36}

Androgens seem to regulate VEGF expression in PCa cells and prostatic fibroblasts.^{37,38} Hypoxia also induces VEGF transcription through the mediator, hypoxia-inducible factor-1,³⁹ whose promoter activity is inhibited by protein kinase C inhibitors.⁴⁰ Therefore, hypoxia, which can occur when tumours outgrow their blood supply, is a likely explanation for the induction of angiogenesis in PCa.

Several human PCa cell lines express enzymes that convert plasminogen or plasmin to the potent angiogenic product angiostatin.^{41,42} Furthermore, Fortier and colleagues found that prostate-specific antigen has dose-dependent antiangiogenic activity and inhibits endothelial cell response to both bFGF and VEGF.⁴³ This antiangiogenic activity may explain the slow progression of certain PCas; however, levels of prostate-specific antigen studied by Fortier and coworkers were, for the most part, much higher than those observed in the serum of patients with PCa. Aberrant nitric oxide production by prostate tumour or host cells may also be a factor in tumour progression because of stimulation mediated by nitric oxide of cancer cell migration and invasiveness.⁴⁴⁻⁴⁶

Although the biology of PCa has been further characterized by the identification of relevant angiogenic factors, additional studies are needed to fully elucidate the multiple mechanisms involved in PCa angiogenesis.

Prognostic markers

Both MVD and the expression of various angiogenic factors have been evaluated as prognostic factors for patients with PCa. MVD is a potential prognostic factor that has been correlated with clinical stage, pathologic stage, metastasis, and histopathologic grade and is a significant predictor of disease-specific survival and progression after therapy.^{23,47-51} However, MVD has not been consistently shown to correlate independently with outcome after radical prostatectomy⁵²⁻⁵⁶ and, therefore, the prognostic value of MVD needs to be confirmed in prospective studies prior to widespread clinical use.

Both VEGF and bFGF have also been evaluated as prognostic factors in PCa, however, neither is prognostic and their detection may merely reflect the presence of benign prostatic hyperplasia, which is usually present in most patients with PCa.^{30,57-59} In preliminary studies, transforming growth factor- β , correlated with tumour grade, MVD, bone metastases and a shorter median cancer-specific survival,⁶⁰ whereas platelet-derived endothelial growth factor correlated with increased MVD.⁶¹

Antiangiogenic therapies

Most studies that have evaluated antiangiogenic therapy for PCa have been in the preclinical setting, and the efficacy demonstrated suggests that these agents may be useful clinically.

Systemic therapy with the anti-VEGF antibody A4.6.1 inhibited neovascularization, the local growth of human PCa xenografts⁶² and lung metastasis.⁶³ Treatment of the PC-3 and Dunning R-3327 cell lines growing subcutaneously in athymic mice with the synthetic fumagillin analogue 0-(chloroacetyl-carbamoyl) fumagillol (TNP-470) inhibited tumour growth and metastasis in a dose-dependent manner.^{63,64} Another antiangiogenic agent, linomide, inhibited tumour-induced angiogenesis and growth of PC-3 tumours implanted subcutaneously in nude mice,^{65,66} and it appeared to act directly on tumours with no significant effect on VEGF or bFGF expression relative to controls.⁶⁷

Interferon- α (IFN- α) and IFN- β are cytokines with antiangiogenic activity. Transfected PCa cells with the murine IFN- β gene may have decreased tumorigenesis and metastasis resulting from bystander effects (the effect of interferon on murine inflammatory cells and stroma), resulting in decreased tumour angiogenesis.⁶⁸ Although interferon has been relatively ineffective in the treatment of most solid tumours clinically, the delivery of optimal doses using effective schedules⁶⁹ and local delivery via gene therapy⁶⁸ may significantly improve the results of interferon therapy.

In a phase II clinical trial, the antiangiogenic agent carboxyamido-triazole was administered to 15 patients with androgen-independent PCa, but no significant clinical response was seen.⁷⁰ PCa angiogenesis is thought to be androgen dependent; however, in a study evaluating neoadjuvant androgen deprivation, no effect on MVD was observed in 80 patients who had undergone radical prostatectomy.⁷¹

Consumption of energy-reduced diets by mice with human PCa xenografts resulted in antiangiogenic activity and decreased VEGF expression,⁷² whereas the consumption of soy products enhanced apoptosis and decreased tumour cell proliferation, MVD and tumour volume.⁷³

Thus far, no reported antiangiogenic therapies have demonstrated significant efficacy in clinical studies. Promising novel and more specific molecules such as the anti-VEGF receptor monoclonal antibody DC101, used as monotherapy or adjuvantly with chemotherapy with or without androgen deprivation, are undergoing preclinical studies.

Bladder cancer

Angiogenesis

The role of angiogenesis in TCC is well established.⁷⁴⁻⁸⁵ Although the expression of bFGF is variable, it appears to be an important mediator of angiogenesis in TCC. Highly metastatic variants of the human 253J TCC cell line expressed high levels of bFGF mRNA compared with the parental cell line and were highly angiogenic and metastatic.⁸⁶ However, the balance between proangiogenic peptides and inhibitors ultimately determines the angiogenic phenotype of the tumour cell, and downregulation of the endogenous angiogenic inhibitor thrombospondin seems to be another key event in bladder carcinogenesis.⁸⁷

Thymidine phosphorylase has angiogenic activity and may also play a role in the pathogenesis of TCC.⁸⁸ Hyaluronidase degrades hyaluronic acid into small fragments that stimulate angiogenesis, and urinary levels of hyaluronic acid, hyaluronidase and hyaluronic acid fragments are elevated in patients with TCC.⁸⁹⁻⁹¹

Prognostic markers

MVD has been extensively evaluated as a prognostic factor for TCC and correlated with lymph node metastasis, disease recurrence and survival, and it stratified the risk of recurrence attributed to p53 expression in muscle-invasive TCC.⁹²⁻⁹⁶ However, MVD was not predictive of disease progression in patients with superficial lamina propria-invasive tumours.⁹⁷ It appears that MVD can be used to identify patients with locally advanced TCC who are at risk of developing metastatic disease after definitive therapy.

VEGF and bFGF expression by human TCC has also been evaluated, and thus far no definitive conclusions can be reached regarding their prognostic value.^{77,98-103} However, bFGF may be a marker of chemoresistance, because the HT1376 TCC cell line, following transfection with the bFGF sense cDNA, demonstrated cisplatin resistance.¹⁰⁴ Overexpression of bFGF mRNA in radical cystectomy specimens also independently predicted poor outcome for patients who received neoadjuvant chemotherapy.⁹⁵

Angiogenin is another proangiogenic factor that is overexpressed in muscle-invasive TCC relative to superficial TCC, and increased serum angiogenin levels were associated with a lower overall survival rate.¹⁰⁵ The loss of expression of thrombospondin protein has also been suggested to be an independent predictor of disease recurrence and overall survival in patients undergoing radical cystectomy.⁸⁵ Invasive tumours have also been found to overexpress the heparin-binding growth factor midkine⁸⁰ and hepatocyte growth factor.^{81,82} Expressions of several angiogenic factors and MVD appear to be potential prognostic markers for TCC, but again prospective studies are necessary before their widespread clinical use.

Antiangiogenic therapies

Several antiangiogenic therapies have demonstrated efficacy for TCC in preclinical studies. Subcutaneous IFN- α therapy inhibited tumour growth and decreased MVD and the expressions of bFGF and VEGF in human TCC growing orthotopically in nude mice.¹⁰⁶ This effect was observed in tumours that were resistant to the antiproliferative effect of interferon, indicating that the inhibition of tumour growth was dependent upon the antiangiogenic effect of interferon. Subsequently, it has been demonstrated that the optimal antiangiogenic effect and efficacy of IFN- α is highly dependent upon the dose and schedule of administration.⁶⁹

Antiangiogenic activity has been observed following therapy for human bladder cancer xenografts with the

antiepidermal growth factor receptor monoclonal antibody C225. Therapy with C225 caused the regression of established human xenografts by downregulating the expression of VEGF, bFGF and IL-8 with concomitant inhibition of growth and metastasis.¹⁰⁷

Similarly, therapy with DC101, in combination with paclitaxel, inhibited tumour growth and metastasis and improved the survival of mice with human bladder cancer xenografts growing orthotopically within the bladder of athymic nude mice.¹⁰⁸ In a heterotopic murine model for bladder TCC, both the pharmaceutical agent halofuginone and dietary soy products have demonstrated antiangiogenic effects.^{109,110}

Renal cell carcinoma

Angiogenesis

RCC is among the most vascular of solid tumours, which suggests a prominent role for neovascularization in its pathogenesis, and experiments have demonstrated that human RCC cells induced angiogenesis.¹¹¹ VEGF may play a prominent role in RCC angiogenesis, because several studies have demonstrated higher expression levels of VEGF mRNA and protein in tumours relative to normal renal tubules¹¹²⁻¹¹⁴ and benign oncocytomas,¹¹⁴ and VEGF appears to act on RCC endothelial cells in a paracrine fashion.^{112,115} VEGF has 5 isoforms and expresses variable levels of each, and the increased expression of VEGF-189 in RCC has been associated with higher stage tumours.¹¹⁶ The role of bFGF in RCC angiogenesis and tumorigenesis is less clear.^{113,117}

The activity of MMPs is regulated by the balance of expression with tissue inhibitors of metalloproteinase, whose inactivation by methylation has been observed in RCC.¹¹⁸ Urokinase-type plasminogen activator (uPA) is a serine protease that activates plasminogen to form plasmin, resulting in alterations of the extracellular matrix degradation, thereby facilitating endothelial cell migration during angiogenesis. Overexpression of the uPA receptor has been observed in RCC.¹¹⁹ Levels of uPA correlate with inactivation of the von Hippel-Lindau gene, a tumour suppressor gene inactivated in the majority (75%) of sporadic RCCs.^{120,121} This suggests that the uPA system plays an important role in angiogenesis, tumorigenicity and metastasis in RCC. The von Hippel-Lindau protein may also be involved in several alternative pathways affecting angiogenesis in RCC by regulating VEGF production and causing regression of the angiogenic mediator transforming growth factor- β_1 .¹²²⁻¹²⁴

Prognostic markers

MVD and the expression of angiogenic factors have been evaluated as prognostic markers for RCC, but it is unclear whether either contributes any additional prognostic information in addition to that obtained from current clini-

copathologic parameters.¹²⁵⁻¹³¹ Thymidine phosphorylase overexpression correlated with MVD and was an independent prognostic factor for survival in RCC.¹²⁵

Angiogenesis factors have also been identified in the serum of patients with RCC. VEGF levels were significantly higher in patients with RCC relative to control patients, but they did not remain a prognostic factor upon multivariate analysis, which included tumour grade and stage.¹³² Associations between bFGF or VEGF levels and tumour grade, stage or overall survival have differed among studies.^{132,133} The VEGF receptor (FLT-1) has also been evaluated as a prognostic factor in RCC, and increased FLT-1 mRNA expression was observed in RCC compared with normal renal tissue.¹³⁴ No angiogenic factor appears to be a candidate prognostic marker for RCC.

Antiangiogenic therapies

Several antiangiogenic therapies for RCC have shown promise in preclinical studies and are currently being evaluated in clinical trials. Systemic administration of the endogenous angiogenic inhibitor endostatin suppressed tumour growth in RCC in a murine model,¹³⁵ and endostatin is now undergoing evaluation in a multi-institutional phase I clinical trial for patients with solid malignant tumours. TNP-470 demonstrated promising results in preclinical studies, however, a phase II clinical study of 33 patients with metastatic RCC found that TNP-470 had manageable toxic effects, but led to only one partial response of short duration.¹³⁶⁻¹³⁸

The results of interferon therapy for locally advanced or metastatic RCC have been disappointing. In a phase II clinical study, subcutaneous IFN- α -2b was used to treat 12 patients with progressive metastatic RCC.¹³⁹ Its efficacy was minimal, although higher responses were observed in patients with higher bFGF serum concentrations. Two well-designed, randomized prospective studies of patients with metastatic RCC did not demonstrate a survival advantage with IFN- α -2a or IFN- γ -1b.^{140,141} Several other phase II or III clinical trials, with less rigorous study designs, have also demonstrated modest responses or no efficacy for interferon therapy with or without chemotherapeutic agents or other immunotherapies in the treatment of metastatic RCC.¹⁴²⁻¹⁵¹ Significant toxicity was experienced in these trials. Only 2 trials have shown that interferon provided a modest survival advantage compared with chemotherapy or hormonal therapy alone.^{152,153} Optimal doses, scheduling and the type of systemic interferon administered may improve the antiangiogenic effect and clinical efficacy.⁶⁸ Vitamin D₃ agents also inhibited angiogenesis, tumorigenicity and metastases in animal xenografts of RCC.¹⁵⁴ Currently, clinical trials are evaluating the efficacy of the angiogenesis inhibitor thalidomide in patients with advanced RCC.

Conclusion

The mechanisms regulating the acquisition of the angio-

genic phenotype of urologic cancers are undergoing vigorous study but have not, as yet, been fully elucidated. Angiogenesis plays a critical role in tumorigenesis and metastasis in prostate, bladder and renal cancers. Several angiogenic biomarkers have been implicated as prognostic markers for urologic malignant diseases in preliminary reports, but these results need to be confirmed in prospective studies. Antiangiogenic therapy targets pathways regulating tumour growth and metastasis, and promising preclinical studies have led to the initiation of phase I trials for patients with muscle-invasive TCC using the antiangiogenic agents C225 with chemotherapy and interferon with chemotherapy in a neoadjuvant setting. These antiangiogenic strategies, especially in combination with cytoreductive chemotherapy, may lead to novel treatments for urologic cancers and may identify new intermediate markers for the response to therapy.

Competing interests: None declared.

Contributors: Both authors contributed to the conception and design of this review and to the interpretation of the data. Both authors contributed equally to the drafting of the manuscript and revising it critically for its scientific content.

Acknowledgement: This review was supported in part by the Royal College of Physicians and Surgeons of Canada Detweiler Traveling Fellowship awarded to Dr. Izawa.

References

- Greenlee RT, Murray T, Bolden S, Wingo PA. Cancer statistics, 2000. *CA Cancer J Clin* 2000;50:7-33.
- Poste G, Fidler IJ. The pathogenesis of cancer metastasis. *Nature* 1979;283:139-46.
- Fidler IJ, Hart IR. Biological diversity in metastatic neoplasms: origins and implications. *Science* 1982;217:998-1003.
- Fidler IJ. Critical factors in the biology of human cancer metastasis: twenty-eighth G.H.A. Clowes Memorial Award Lecture. *Cancer Res* 1990;50:6130-8.
- Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000;100:57-70.
- Folkman J. How is blood vessel growth regulated in normal and neoplastic tissue? G.H.A. Clowes Memorial Award Lecture. *Cancer Res* 1986;46:467-73.
- Folkman J, Watson K, Ingber D, Hanahan D. Induction of angiogenesis during the transition from hyperplasia to neoplasia. *Nature* 1989;339:58-61.
- Liotta LA, Steeg PS, Stetler-Stevenson WG. Cancer metastasis and angiogenesis: an imbalance of positive and negative regulation. *Cell* 1991;64:327-36.
- Fidler IJ, Ellis LM. The implications of angiogenesis for the biology and therapy of cancer metastasis. *Cell* 1994;79:185-8.
- Folkman J. Clinical applications of research on angiogenesis. *N Engl J Med* 1995;333:1757-63.
- Folkman J. Anti-angiogenesis: new concept for therapy of solid tumors. *Ann Surg* 1972;175:409-16.
- Gimbrone MA Jr, Cotran RS, Leapman SB, Folkman J. Tumor growth and neovascularization: an experimental model using the rabbit cornea. *J Natl Cancer Inst* 1974;52:413-27.
- Hanahan D, Folkman J. Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* 1996;86:353-64.
- Folkman J. Angiogenesis in cancer, vascular, rheumatoid and other disease. *Nat Med* 1995;1:27-31.
- Singh RK, Llansa N, Bucana CD, Sanchez R, Koura A, Fidler IJ. Cell density-dependent regulation of basic fibroblast growth factor expression in human renal cell carcinoma cells. *Cell Growth Differ* 1996;7:397-404.
- Naito S, von Eschenbach AC, Giavazzi R, Fidler IJ. Growth and metastasis of tumor cells isolated from a human renal cell carcinoma implanted into different organs of nude mice. *Cancer Res* 1986;46:4109-15.
- Fidler IJ, Naito S, Pathak S. Orthotopic implantation is essential for the selection, growth and metastasis of human renal cell cancer in nude mice. *Cancer Metastasis Rev* 1990;9:149-65.
- Fidler IJ, Wilmanns C, Staroselsky A, Radinsky R, Dong Z, Fan D. Modulation of tumor cell response to chemotherapy by the organ environment. *Cancer Metastasis Rev* 1994;13:201-22.
- Perrotte P, Bielenberg DR, Eve BY, Dinney CPN. Organ-specific angio-

- genesis and metastases of human bladder carcinoma growing in athymic nude mice. *Mol Urol* 1997;1:299-307.
20. Staroselsky AN, Radinsky R, Fidler IJ, Pathak S, Chernajovsky Y, Frost P. The use of molecular genetic markers to demonstrate the effect of organ environment on clonal dominance in a human renal-cell carcinoma grown in nude mice. *Int J Cancer* 1992;51:130-8.
 21. Hanahan D, Christofori G, Naik P, Arbeit J. Transgenic mouse models of tumour angiogenesis: the angiogenic switch, its molecular controls, and prospects for preclinical therapeutic models. *Eur J Cancer* 1996;32A:2386-93.
 22. Kumar R, Fidler IJ. Angiogenic molecules and cancer metastasis. *In Vivo* 1998;18:27-34.
 23. Weidner N, Carroll PR, Flax J, Blumenfeld W, Folkman J. Tumor angiogenesis correlates with metastasis in invasive prostate carcinoma. *Am J Pathol* 1993;143:401-9.
 24. Hayes AJ, Li LY, Lippman ME. Science, medicine, and the future. Antivascular therapy: a new approach to cancer treatment. *BMJ* 1999;318:853-6.
 25. Brooks PC, Montgomery AM, Rosenfeld M, Reisfeld RA, Hu T, Klier G, et al. Integrin alpha v beta 3 antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell* 1994;79:1157-64.
 26. Huang X, Molema G, King S, Watkins L, Edgington TS, Thorpe PE. Tumor infarction in mice by antibody-directed targeting of tissue factor to tumor vasculature. *Science* 1997;275:547-50.
 27. Boehm T, Folkman J, Browder T, O'Reilly MS. Antiangiogenic therapy of experimental cancer does not induce acquired drug resistance. *Nature* 1997;390:404-7.
 28. Holmgren L, O'Reilly MS, Folkman J. Dormancy of micrometastases: balanced proliferation and apoptosis in the presence of angiogenesis suppression. *Nat Med* 1995;1:149-53.
 29. Hepburn PJ, Griffiths K, Harper ME. Angiogenic factors expressed by human prostatic cell lines: effect on endothelial cell growth in vitro. *Prostate* 1997;33:123-32.
 30. Weingartner K, Ben-Sasson SA, Stewart R, Richie JP, Riedmiller H, Folkman J. Endothelial cell proliferation activity in benign prostatic hyperplasia and prostate cancer: an in vitro model for assessment. *J Urol* 1998;159:465-70.
 31. Ferrer FA, Miller LJ, Andrawis RI, Kurtzman SH, Albertson PC, Laudone VP, et al. Angiogenesis and prostate cancer: in vivo and in vitro expression of angiogenesis factors by prostate cancer cells. *Urology* 1998;51:161-7.
 32. Greene GF, Kitadai Y, Pettaway CA, von Eschenbach AC, Bucana CD, Fidler IJ. Correlation of metastasis-related gene expression with metastatic potential in human prostate carcinoma cells implanted in nude mice using an in situ messenger RNA hybridization technique. *Am J Pathol* 1997;150:1571-82.
 33. Balbay MD, Pettaway CA, Kuniyasu H, Inoue K, Ramirez E, Li E, et al. Highly metastatic human prostate cancer growing within the prostate of athymic mice overexpresses vascular endothelial growth factor. *Clin Cancer Res* 1999;5:783-9.
 34. Inoue K, Slaton JW, Eve BY, Kim SJ, Perrotte P, Balbay D, et al. Interleukin-8 expression regulates tumorigenicity and metastases in androgen-independent prostate cancer. *Clin Cancer Res* 2000;6:2104-19.
 35. Stearns ME, Garcia FU, Fudge K, Rhim J, Wang M. Role of interleukin 10 and transforming growth factor beta1 in the angiogenesis and metastasis of human prostate primary tumor lines from orthotopic implants in severe combined immunodeficiency mice. *Clin Cancer Res* 1999;5:711-20.
 36. Stearns ME, Rhim J, Wang M. Interleukin 10 (IL-10) inhibition of primary human prostate cell-induced angiogenesis: IL-10 stimulation of tissue inhibitor of metalloproteinase-1 and inhibition of matrix metalloproteinase (MMP)-2/MMP-9 secretion. *Clin Cancer Res* 1999;5:189-96.
 37. Joseph IB, Nelson JB, Denmeade SR, Isaacs JT. Androgens regulate vascular endothelial growth factor content in normal and malignant prostatic tissue. *Clin Cancer Res* 1997;3:2507-11.
 38. Levine AC, Liu XH, Greenberg PD, Eliashvili M, Schiff JD, Aaronson SA, et al. Androgens induce the expression of vascular endothelial growth factor in human fetal prostatic fibroblasts. *Endocrinology* 1998;139:4672-8.
 39. Zhong H, De Marzo AM, Laughner E, Lim M, Hilton DA, Zagzag D, et al. Overexpression of hypoxia-inducible factor 1alpha in common human cancers and their metastases. *Cancer Res* 1999;59:5830-5.
 40. Kruger EA, Blagosklonny MV, Dixon SC, Figg WD. UCN-01, a protein kinase C inhibitor, inhibits endothelial cell proliferation and angiogenic hypoxic response. *Invasion Metastasis* 2000;18:209-18.
 41. Gately S, Twardowski P, Stack MS, Patrick M, Boggio L, Cundiff DL, et al. Human prostate carcinoma cells express enzymatic activity that converts human plasminogen to the angiogenesis inhibitor, angiotatin. *Cancer Res* 1996;56:4887-90.
 42. Heidtmann HH, Nettelbeck DM, Mingels A, Jager R, Welker HG, Kontermann RE. Generation of angiotatin-like fragments from plasminogen by prostate-specific antigen. *Br J Cancer* 1999;81:1269-73.
 43. Fortier AH, Nelson BJ, Grella DK, Holaday JW. Antiangiogenic activity of prostate-specific antigen. *J Natl Cancer Inst* 1999;91:1635-40.
 44. Thomsen LL, Miles DW. Role of nitric oxide in tumour progression: lessons from human tumours. *Cancer Metastasis Rev* 1998;17:107-18.
 45. Jadeski LC, Lala PK. Nitric oxide synthase inhibition by N(G)-nitro-L-arginine methyl ester inhibits tumor-induced angiogenesis in mammary tumors. *Am J Pathol* 1999;155:1381-90.
 46. Klotz T, Bloch W, Volberg C, Engelmann U, Addicks K. Selective expression of inducible nitric oxide synthase in human prostate carcinoma. *Cancer* 1998;82:1897-1903.
 47. Fregene TA, Khanuja PS, Noto AC, Gehani SK, van Egmont EM, Luz DA, et al. Tumor-associated angiogenesis in prostate cancer. *Anticancer Res* 1993;13:2377-82.
 48. Borre M, Offersen BV, Nerstrom B, Overgaard J. Microvessel density predicts survival in prostate cancer patients subjected to watchful waiting. *Br J Cancer* 1998;78:940-4.
 49. Brauer MK, Deering RE, Brown M, Preston SD, Bigler SA. Predictors of pathologic stage in prostatic carcinoma. The role of neovascularity. *Cancer* 1994;73:678-87.
 50. Silberman MA, Partin AW, Veltri RW, Epstein JI. Tumor angiogenesis correlates with progression after radical prostatectomy but not with pathologic stage in Gleason sum 5 to 7 adenocarcinoma of the prostate. *Cancer* 1997;79:772-9.
 51. Strohmeier D, Rossing C, Strauss F, Bauerfeind A, Kaufmann O, Loening S. Tumor angiogenesis is associated with progression after radical prostatectomy in pT2/pT3 prostate cancer. *Prostate* 2000;42:26-33.
 52. Hall MC, Troncoso P, Pollack A, Zhou HYE, Zagars GK, Chung LWK, et al. Significance of tumor angiogenesis in clinically localized prostatic carcinoma treated with external beam radiotherapy. *Urology* 1994;44:869-75.
 53. Bettencourt MC, Bauer JJ, Sesterhenn IA, Connelly RR, Moul JW. CD34 immunohistochemistry assessment of angiogenesis as a prognostic marker for prostate cancer recurrence after radical prostatectomy. *J Urol* 1998;160:459-65.
 54. Moul JW. Angiogenesis, p53, bcl-2 and Ki-67 in the progression of prostate cancer after radical prostatectomy. *Eur Urol* 1999;35:399-407.
 55. Rubin MA, Buyyounouski M, Bagiella E, Sharir S, Neugut A, Benson M, et al. Microvessel density in prostate cancer: lack of correlation with tumor grade, pathologic stage, and clinical outcome. *Urology* 1999;53:542-7.
 56. Gettman MT, Parelli A, Slezak J, Bergstralh EJ, Blute M, Zincke H, et al. Role of microvessel density in predicting recurrence in pathologic stage T3 prostatic adenocarcinoma. *Urology* 1999;54:479-85.
 57. Duque JL, Loughlin KR, Adam RM, Kantoff PW, Zurakowski D, Freeman MR. Plasma levels of vascular endothelial growth factor are increased in patients with metastatic prostate cancer. *Urology* 1999;54:523-7.
 58. Walsh K, Sherwood RA, Dew TK, Mulvin D. Angiogenic peptides in prostate disease. *Br J Urol* 1999;84:1081-3.
 59. Meyer GE, Yu E, Siegal JA, Petteway JC, Blumenstein BA, Brawer MK. Serum basic fibroblast growth factor in men with and without prostate carcinoma. *Cancer* 1995;76:2304-11.
 60. Wikstrom P, Stattin P, Franck-Lissbrant I, Damber JE, Bergh A. Transforming growth factor beta1 is associated with angiogenesis, metastasis, and poor clinical outcome in prostate cancer. *Prostate* 1998;37:19-29.
 61. Sugamoto T, Tanji N, Nishio S, Yokoyama M. Expression of platelet-derived endothelial cell growth factor in prostatic adenocarcinoma. *Oncol Rep* 1999;6:519-22.
 62. Borgstrom P, Bourdon MA, Hillan KJ, Sriramarao P, Ferrara N. Neutralizing anti-vascular endothelial growth factor antibody completely inhibits angiogenesis and growth of human prostate carcinoma micro tumors in vivo. *Prostate* 1998;35:1-10.
 63. Miki T, Nonomura N, Nozawa M, Harada Y, Nishimura K, Kojima Y, et al. Angiogenesis inhibitor TNP-470 inhibits growth and metastasis of a hormone-independent rat prostatic carcinoma cell line. *J Urol* 1998;160:210-3.
 64. Yamaoka M, Yamamoto T, Ikeyama S, Sudo K, Fujita T. Angiogenesis inhibitor TNP-470 (AGM-1470) potently inhibits the tumor growth of hormone-independent human breast and prostate carcinoma cell lines. *Cancer Res* 1993;53:5233-6.
 65. Vukanovic J, Isaacs JT. Human prostatic cancer cells are sensitive to programmed (apoptotic) death induced by the antiangiogenic agent linomide. *Cancer Res* 1995;55:3517-20.
 66. Joseph IB, Isaacs JT. Potentiation of the antiangiogenic ability of linomide by androgen ablation involves down-regulation of vascular endothelial growth factor in human androgen-responsive prostatic carcinomas. *Cancer Res* 1997;57:1054-7.
 67. Joseph IB, Isaacs JT. Macrophage role in the anti-prostate cancer response to one class of antiangiogenic agents. *J Natl Cancer Inst* 1998;90:1648-53.
 68. Dong Z, Greene G, Pettaway C, Dinney CPN, Eue I, Lu W, et al. Suppression of angiogenesis, tumorigenicity, and metastasis by human prostate cancer cells engineered to produce interferon-beta. *Cancer Res* 1999;59:872-9.
 69. Slaton JW, Perrotte P, Inoue K, Dinney CPN, Fidler IJ. Interferon-alpha-mediated down-regulation of angiogenesis-related genes and therapy of bladder cancer are dependent on optimization of biological dose and schedule. *Clin Cancer Res* 1999;5:2726-34.
 70. Bauer KS, Figg WD, Hamilton JM, Jones EC, Premkumar A, Steinberg SM, et al. A pharmacokinetically guided phase II study of carboxyamido-triazole in androgen-independent prostate cancer. *Clin Cancer Res* 1999;5:2324-9.
 71. Matsushima H, Goto T, Hosaka Y, Kitamura T, Kawabe K. Correlation between proliferation, apoptosis, and angiogenesis in prostate carcinoma and their relation to androgen ablation. *Cancer* 1999;85:1822-7.
 72. Mukherjee P, Sotnikov AV, Mangian HJ, Zhou J, Visek WJ, Clinton SK. En-

- ergy intake and prostate tumor growth, angiogenesis, and vascular endothelial growth factor expression. *J Natl Cancer Inst* 1999;91:512-23.
73. Zhou J, Gugger ET, Tanaka T, Guo Y, Blackburn GL, Clinton SK. Soybean phytochemicals inhibit the growth of transplantable human prostate carcinoma and tumor angiogenesis in mice. *J Nutr* 1999;129:1628-35.
 74. Chodak GW, Scheiner CJ, Zetter BR. Urine from patients with transitional-cell carcinoma stimulates migration of capillary endothelial cells. *N Engl J Med* 1981;305:869-74.
 75. Chodak GW, Summerhayes I. Detection of angiogenesis activity in malignant bladder tissue and cells. *J Urol* 1984;132:1032-5.
 76. Chopin DK, Caruelle JP, Columbel M, Palcy S, Ravery V, Caruelle D, et al. Increased immunodetection of acidic fibroblast growth factor in bladder cancer, detectable in urine. *J Urol* 1993;150:1126-30.
 77. Nguyen M, Watanabe H, Budson AE, Richie JP, Folkman J. Elevated levels of the angiogenic peptide basic fibroblast growth factor in urine of bladder cancer patients. *J Natl Cancer Inst* 1993;85:241-2.
 78. O'Brien T, Cranston D, Fuggle S, Bicknell R, Harris AL. Two mechanisms of basic fibroblast growth factor-induced angiogenesis in bladder cancer. *Cancer Res* 1997;57:136-40.
 79. O'Brien T, Cranston D, Fuggle S, Bicknell R, Harris AL. Different angiogenic pathways characterize superficial and invasive bladder cancer. *Cancer Res* 1995;55:510-3.
 80. O'Brien TS, Cranston D, Fuggle S, Bicknell R, Harris AL. The angiogenic factor midkine is expressed in bladder cancer, and overexpression correlates with a poor outcome in patients with invasive cancers. *Cancer Res* 1996;56:2515-8.
 81. Joseph A, Weiss GH, Jin L, Fuchs A, Chowdhury S, O'Shaughnessy P, et al. Expression of scatter factor in human bladder carcinoma. *J Natl Cancer Inst* 1995;87:372-7.
 82. Rosen EM, Joseph A, Jin L, Yao Y, Chau MH, Fuchs A, et al. Urinary and tissue levels of scatter factor in transitional cell carcinoma of bladder. *J Urol* 1997;157:72-8.
 83. Tachibana M, Miyakawa A, Nakashima J, Murai M, Nakamura K, Kubo A, et al. Constitutive production of multiple cytokines and a human chorionic gonadotropin beta-subunit by a human bladder cancer cell line (KU-19-19): possible demonstration of totipotential differentiation. *Br J Cancer* 1997;76:163-74.
 84. Eder IE, Stenzl A, Hobisch A, Cronauer MV, Bartsch G, Klocker H. Expression of TGF β 's beta-1, beta 2 and beta 3 in human bladder carcinomas. *Br J Cancer* 1997;75:1753-60.
 85. Grossfeld GD, Ginsberg DA, Stein JP, Bochner BH, Esrig D, Groshen S, et al. Thrombospondin-1 expression in bladder cancer: association with p53 alterations, tumor angiogenesis, and tumor progression. *J Natl Cancer Inst* 1997;89:219-27.
 86. Dinney CP, Fishbeck R, Singh RK, Eve B, Pathak S, Brown N, et al. Isolation and characterization of metastatic variants from human transitional cell carcinoma passaged by orthotopic implantation in athymic nude mice. *J Urol* 1995;154:1532-8.
 87. Campbell SC, Volpert OV, Ivanovich M, Bouck NP. Molecular mediators of angiogenesis in bladder cancer. *Cancer Res* 1998;58:1298-1304.
 88. Kubota Y, Miura T, Moriyama M, Noguchi S, Matsuzaki J, Takebayashi S, et al. Thymidine phosphorylase activity in human bladder cancer: difference between superficial and invasive cancer. *Clin Cancer Res* 1997;3:973-6.
 89. Lokeshwar VB, Obek C, Soloway MS, Block NL. Tumor-associated hyaluronic acid: a new sensitive and specific urine marker for bladder cancer. *Cancer Res* 1997;57:773-7.
 90. Lokeshwar VB, Obek C, Pham HT, Wei D, Young MJ, Duncan RC, et al. Urinary hyaluronic acid and hyaluronidase: markers for bladder cancer detection and evaluation of grade. *J Urol* 2000;163:348-56.
 91. Lokeshwar VB, Young MJ, Goudarzi G, Iida N, Yudin AI, Cherr GN, et al. Identification of bladder tumor-derived hyaluronidase: its similarity to HYAL1. *Cancer Res* 1999;59:4464-70.
 92. Dickenson AJ, Fox SB, Persad RA, Hollyer J, Sibley GNA, Harris AL. Quantification of angiogenesis as an independent predictor of prognosis in invasive bladder carcinomas. *Br J Urol* 1994;74:762-6.
 93. Bochner BH, Cote RJ, Weidner N, Groshen S, Chen S, Skinner DG, et al. Angiogenesis in bladder cancer: relationship between microvessel density and tumor prognosis. *J Natl Cancer Inst* 1995;87:1603-12.
 94. Jaeger TM, Weidner N, Chew K, Moore DH, Kerschmann RL, Waldman FM, et al. Tumor angiogenesis correlates with lymph node metastases in invasive bladder cancer. *J Urol* 1995;154:69-71.
 95. Chaudhary R, Bromley M, Clarke NW, Betts CD, Barnard RJ, Ryder WD, et al. Prognostic relevance of micro-vessel density in cancer of the urinary bladder. *Anticancer Res* 1999;19:3479-84.
 96. Bochner BH, Esrig D, Groshen S, Dickinson M, Weidner N, Nichols PW, et al. Relationship of tumor angiogenesis and nuclear p53 accumulation in invasive bladder cancer. *Clin Cancer Res* 1997;3:1615-22.
 97. Dinney CP, Babkowski RC, Antelo M, Perrotte P, Liebert M, Zhang HZ, et al. Relationship among cystectomy, microvessel density and prognosis in stage T1 transitional cell carcinoma of the bladder. *J Urol* 1998;160:1285-90.
 98. Gazzaniga P, Gandini O, Gradilone A, Silvestri I, Giuliani L, Magnanti M, et al. Detection of basic fibroblast growth factor mRNA in urinary bladder cancer: correlation with local relapses. *Int J Oncol* 1999;14:1123-7.
 99. Crew JP, O'Brien T, Bicknell R, Fuggle S, Cranston D, Harris AL. Urinary vascular endothelial growth factor and its correlation with bladder cancer recurrence rates. *J Urol* 1999;161:799-804.
 100. Edgren M, Lennernas B, Larsson A, Nilsson S. Serum concentrations of VEGF and b-FGF in renal cell, prostate and urinary bladder carcinomas. *Anticancer Res* 1999;19:869-74.
 101. Crew JP, O'Brien T, Bradburn M, Fuggle S, Bicknell R, Cranston D, et al. Vascular endothelial growth factor is a predictor of relapse and stage progression in superficial bladder cancer. *Cancer Res* 1997;57:5281-5.
 102. Crew JP, Fuggle S, Bicknell R, Cranston DW, de Benedetti A, Harris AL. Eukaryotic initiation factor-4E in superficial and muscle invasive bladder cancer and its correlation with vascular endothelial growth factor expression and tumour progression. *Br J Cancer* 2000;82:161-6.
 103. Chow NH, Liu HS, Chan SH, Cheng HL, Tzai TS. Expression of vascular endothelial growth factor in primary superficial bladder cancer. *Anticancer Res* 1999;19:4593-7.
 104. Miyake H, Hara I, Gohji K, Yoshimura K, Arakawa S, Kamidono S. Expression of basic fibroblast growth factor is associated with resistance to cisplatin in a human bladder cancer cell line. *Cancer Lett* 1998;123:121-6.
 105. Miyake H, Hara I, Yamanaka K, Gohji K, Arakawa S, Kamidono S. Increased angiogenin expression in the tumor tissue and serum of urothelial carcinoma patients is related to disease progression and recurrence. *Cancer* 1999;86:316-24.
 106. Dinney CP, Bielenberg DR, Perrotte P, Reich R, Eve BY, Bucana CD, et al. Inhibition of basic fibroblast growth factor expression, angiogenesis, and growth of human bladder carcinoma in mice by systemic interferon- α administration. *Cancer Res* 1998;58:808-14.
 107. Perrotte P, Matsumoto T, Inoue K, Kuniyasu H, Eve BY, Hicklin DJ, et al. Anti-epidermal growth factor receptor antibody C225 inhibits angiogenesis in human transitional cell carcinoma growing orthotopically in nude mice. *Clin Cancer Res* 1999;5:257-64.
 108. Inoue K, Slaton JW, Davis DW, Hicklin DJ, McConkey DJ, et al. Treatment of human metastatic transitional cell carcinoma of the bladder in a murine model with the anti-vascular endothelial growth factor receptor monoclonal antibody DC101 and paclitaxel. *Clin Cancer Res* 2000;6(7):2635-43.
 109. Elkin M, Ariel I, Miao H, Nagler A, Pines M, de-Groot N, et al. Inhibition of bladder carcinoma angiogenesis, stromal support, and tumor growth by halofuginone. *Cancer Res* 1999;59:4111-8.
 110. Zhou J, Mukherjee P, Gugger ET, Tanaka T, Blackburn GL, Clinton SK. Inhibition of murine bladder tumorigenesis by soy isoflavones via alterations in the cell cycle, apoptosis, and angiogenesis. *Cancer Res* 1998;58:5231-8.
 111. Bard RH, Mydlo JH, Freed SZ. Detection of tumor angiogenesis factor in adenocarcinoma of kidney. *Urology* 1986;27:447-50.
 112. Brown LF, Berse B, Jackman RW, Tognazzi K, Manseau EJ, Dvorak HF, et al. Increased expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in kidney and bladder carcinomas. *Am J Pathol* 1993;143:1255-62.
 113. Takahashi A, Sasaki H, Kim SJ, Tobisu K, Kakizoe T, Tsukamoto T, et al. Markedly increased amounts of messenger RNAs for vascular endothelial growth factor and placenta growth factor in renal cell carcinoma associated with angiogenesis. *Cancer Res* 1994;54:4233-7.
 114. Nicol D, Hii S, Walsh M, The B, Thompson L, Kennett C, et al. Vascular endothelial growth factor expression is increased in renal cell carcinoma. *J Urol* 1997;157:1482-6.
 115. Nakagawa M, Emoto A, Hanada T, Nasu N, Nomura Y. Tubulogenesis by microvascular endothelial cells is mediated by vascular endothelial growth factor (VEGF) in renal cell carcinoma. *Br J Urol* 1997;79:681-7.
 116. Tomisawa M, Tokunaga T, Oshika Y, Tsuchida T, Fukushima Y, Sato H, et al. Expression pattern of vascular endothelial growth factor isoform is closely correlated with tumour stage and vascularisation in renal cell carcinoma. *Eur J Cancer* 1999;35:133-7.
 117. Emoto A, Nakagawa M, Wakabayashi Y, Hanada T, Naito S, Nomura Y. Induction of tubulogenesis of microvascular endothelial cells by basic fibroblast growth factor from human SN12C renal cancer cells. *J Urol* 1997;157:699-703.
 118. Bachman KE, Herman JG, Corn PG, Merlo A, Costello JF, Cavenee WK, et al. Methylation-associated silencing of the tissue inhibitor of metalloproteinase-3 gene suggests a suppressor role in kidney, brain, and other human cancers. *Cancer Res* 1999;59:798-802.
 119. Xu Y, Hagege J, Doublet J, Callard P, Sraer J, Ronne E, et al. Endothelial and macrophage upregulation of urokinase receptor expression in human renal cell carcinoma. *Hum Pathol* 1997;28:206-13.
 120. Foster K, Prowse A, van den Berg A, Fleming S, Hulsbeek MM, Crossey PA, et al. Somatic mutations of the von-Hippel-Lindau disease tumour suppressor gene in non-familial clear cell renal carcinoma. *Hum Mol Genet* 1994;3:2169-73.
 121. Gnarr JR, Tory K, Weng Y, Schmidt L, Wei MH, Li H, et al. Mutations of the VHL tumour suppressor gene in renal carcinoma. *Nat Genet* 1994;7:85-90.
 122. Siemester G, Weindel K, Mohrs K, Barleon B, Martiny-Baron G, Marme D. Reversion of deregulated expression of vascular endothelial growth factor in human renal cell carcinoma cells by von Hippel-Lindau tumor suppressor protein. *Cancer Res* 1996;56:2299-301.

123. Pal S, Claffey KP, Dvorak HF, Mukhopadhyay D. The von Hippel-Lindau gene product inhibits vascular permeability factor/vascular endothelial growth factor expression in renal cell carcinoma by blocking protein kinase C pathways. *J Biol Chem* 1997;272:27509-12.
124. Ananth S, Knebelmann B, Gruning W, Dhanabal M, Walz G, Stillman IE, et al. Transforming growth factor β 1 is a target for the von Hippel-Lindau tumor suppressor and a critical growth factor for clear cell renal carcinoma. *Cancer Res* 1999;59:2210-6.
125. Imazano Y, Takebayashi Y, Nishiyama K, Akiba S, Miyadera K, Yamada Y, et al. Correlation between thymidine phosphorylase expression and prognosis in human renal cell carcinoma. *J Clin Oncol* 1997;15:2570-8.
126. MacLennan GT, Bostwick DG. Microvessel density in renal cell carcinoma: lack of prognostic significance. *Urology* 1995;46:27-30.
127. Gelb AB, Sudilovsky D, Wu CD, Weiss LM, Medeiros LJ. Appraisal of intratumoral microvessel density, MIB-1 score, DNA content, and p53 protein expression as prognostic indicators in patients with locally confined renal cell carcinoma. *Cancer* 1997;80:1768-75.
128. Herbst C, Kosmehl H, Stiller KJ, Berndt A, Eiselt M, Schubert J, et al. Evaluation of microvessel density by computerized image analysis in human renal cell carcinoma. Correlation to pT category, nuclear grade, proliferative activity and occurrence of metastasis. *J Cancer Res Clin Oncol* 1998;124:141-7.
129. Yoshino S, Kato M, Okada K. Prognostic significance of microvessel count in low stage renal cell carcinoma. *Int J Urol* 1995;2:156-60.
130. Nativ O, Sabo E, Reiss A, Wald M, Madjar S, Moskovitz B. Clinical significance of tumor angiogenesis in patients with localized renal cell carcinoma. *Urology* 1998;51:693-6.
131. Delahunt B, Bethwaite PB, Thornton A. Prognostic significance of microscopic vascularity for clear cell renal cell carcinoma. *Br J Urol* 1997;80:401-4.
132. Dosquet C, Coudert M, Lepage E, Cabane J, Richard F. Are angiogenic factors, cytokines, and soluble adhesion molecules prognostic factors in patients with renal cell carcinoma? *Clin Cancer Res* 1997;3:2451-8.
133. Wechsler HW, Bichler K, Feil G, Loeser W, Lahme S, Petri E. Renal cell carcinoma: relevance of angiogenic factors. *Anticancer Res* 1999;19:1537-40.
134. Takahashi A, Sasaki H, Kim SJ, Kakizoe T, Miyao N, Sugimura T, et al. Identification of receptor genes in renal cell carcinoma associated with angiogenesis by differential hybridization technique. *Biochem Biophys Res Commun* 1999;257:855-9.
135. Dhanabal M, Ramchandran R, Volk R, Stillman IE, Lombardo M, Iruela-Arispe ML, et al. Endostatin: yeast production, mutants, and antitumor effect in renal cell carcinoma. *Cancer Res* 1999;59:189-97.
136. Morita T, Shinohara N, Tokue A. Antitumor effect of a synthetic analogue of fumagillin on murine renal carcinoma. *Br J Urol* 1994;74:416-21.
137. Fujioka T, Hasegawa M, Ogiu K, Matsushita Y, Sato M, Kubo T. Antitumor effects of angiogenesis inhibitor 0-(choroacetyl-carbamoyl) fumagillol (TNP-470) against murine renal cell carcinoma. *J Urol* 1996;155:1775-8.
138. Stadler WM, Kuzel T, Shapiro C, Sosman J, Clark J, Vogelzang NJ. Multi-institutional study of the angiogenesis inhibitor TNP-470 in metastatic renal carcinoma. *J Clin Oncol* 1999;17:2541-5.
139. Vermeulen PB, Dirix LY, Martin M, Lemmens J, Van Oosterom AT. Serum basic fibroblast growth factor and vascular endothelial growth factor in metastatic renal cell carcinoma treated with interferon alfa-2b. *J Natl Cancer Inst* 1997;89:1316-7.
140. Negrier S, Escudier B, Lasset C, Douillard JY, Savary J, Chevreau C, et al. Recombinant human interleukin-2, recombinant human interferon alfa-2a, or both in metastatic renal-cell carcinoma. *N Engl J Med* 1998;338:1272-8.
141. Gleave ME, Elhilali M, Fradet Y, Davis I, Venner P, Saad F, et al. Interferon gamma-1b compared with placebo in metastatic renal-cell carcinoma. Canadian Urologic Oncology Group. *N Engl J Med* 1998;338:1265-71.
142. Henriksson R, Nilsson S, Colleen S, Wersall P, Helsing M, Zimmerman R, et al. Survival in renal cell carcinoma—a randomized evaluation of tamoxifen vs interleukin 2, alpha-interferon (leucocyte) and tamoxifen. *Br J Cancer* 1998;77:1311-7.
143. Stadler WM, Kuzel T, Dumas M, Vogelzang NJ. Multicenter phase II trial of interleukin-2, interferon-alpha, and 13-cis-retinoic acid in patients with metastatic renal-cell carcinoma. *J Clin Oncol* 1998;16:1820-5.
144. Tourani JM, Pfister C, Berdah JF, Benhammouda A, Salze P, Monnier A, et al. Outpatient treatment with subcutaneous interleukin-2 and interferon alfa administration in combination with fluorouracil in patients with metastatic renal cell carcinoma: results of a sequential nonrandomized phase II study. Subcutaneous Administration Propeukin Program Cooperative Group. *J Clin Oncol* 1998;16:2505-13.
145. Jayson GC, Middleton M, Lee SM, Ashcroft L, Thatcher N. A randomized phase II trial of interleukin 2 and interleukin 2-interferon alpha in advanced renal cancer. *Br J Cancer* 1998;78:366-9.
146. Ravaud A, Audhuy B, Gomez F, Escudier B, Lesimple T, Chevreau C, et al. Subcutaneous interleukin-2, interferon alfa-2a, and continuous infusion of fluorouracil in metastatic renal cell carcinoma: a multicenter phase II trial. Groupe Francais d'Immunotherapie. *J Clin Oncol* 1998;16:2728-32.
147. Igarashi T, Marumo K, Onishi T, Kobayashi M, Aiba K, Tsushima T, et al. Interferon-alpha and 5-fluorouracil therapy in patients with metastatic renal cell cancer: an open multicenter trial. The Japanese Study Group Against Renal Cancer. *Urology* 1999;53:53-9.
148. Clark JI, Gaynor ER, Martone B, Budds SC, Manjunath R, Flanigan RC, et al. Daily subcutaneous ultra-low-dose interferon-alpha in patients with advanced renal cell carcinoma. *Clin Cancer Res* 1999;5:2374-80.
149. Soori GS, Schulof RS, Stark JJ, Wiemann MC, Honeycutt PJ, Church CK, et al. Continuous-infusion floxuridine and alpha interferon in metastatic renal cancer: a national biotherapy study group phase II study. *Cancer Invest* 1999;17:379-84.
150. Rohde D, Thiemann D, Wildberger J, Wolff J, Jakse G. Treatment of renal cancer patients with gemcitabine (2',2'-difluorodeoxycytidine) and interferons: antitumor activity and toxicity. *Oncol Rep* 1998;5:1555-1560.
151. Fujii A, Yui-En K, Ono Y, Yamamoto H, Gohji K, Takenaka A. Preliminary results of the alternating administration of natural interferon-alpha and recombinant interferon-gamma for metastatic renal cell carcinoma. *Br J Urol* 1999;84:399-404.
152. Pyrhonen S, Salminen E, Ruutu M, Lehtonen T, Nurmi M, Tammela T, et al. Prospective randomized trial of interferon alfa-2a plus vinblastine versus vinblastine alone in patients with advanced renal cell cancer. *J Clin Oncol* 1999;17:2859-67.
153. Medical Research Council Renal Cancer Collaborators. Interferon-alpha and survival in metastatic renal carcinoma: early results of a randomised controlled trial. *Lancet* 1999;353:14-7.
154. Fujioka T, Hasegawa M, Ishikura K, Matsushita Y, Sato M, Tanji S. Inhibition of tumor growth and angiogenesis by vitamin D₃ agents in murine renal cell carcinoma. *J Urol* 1998;160:247-51.

Reprint requests to: Dr. Colin P.N. Dinney, Associate Professor of Urology and Cancer Biology, Department of Urology, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., PO Box 110, Houston TX 77030; fax 713 794-4824; cdinney@notes.mdacc.tmc.edu

Consult The Cochrane Library

This electronic library provides evidence for informed health care decision-making in 6 databases:



- The Cochrane Database of Systematic Reviews
- The Database of Abstracts of Reviews of Effectiveness
- The Health Technology Assessment Database

\$323.95/CMA members, \$359.95/nonmembers. All orders must be prepaid. Please add 7% GST/HST (as applicable), 8% PST (Ontario residents only) and \$4 shipping/handling. Please contact CMA Member Service Centre for network prices. Available on CD-ROM or the Internet.

- The Cochrane Controlled Trials Register
- The Cochrane Review Methodology Database
- The NHS Economic Evaluation Database

CMA Member Service Centre
tel 888 855-2555 or 613 731-8610 x2307
cmamsc@cma.ca

