

Reference

1. Schünemann HJ, Best D, Vist G, Oxman AD, for the GRADE Working Group. Letters, numbers, symbols and words: how to communicate grades of evidence and recommendations [editorial]. *CMAJ* 2003;169(7):677-80.

DOI:10.1053/cmaj.1040014

SARS and the removal of personal protective equipment

In a reported cluster of cases of SARS among Canadian health care workers,¹ infections occurred despite apparent compliance with recommended infection control precautions. In the report it was noted that contact with a patient or with a contaminated environment might have led to health care workers infecting themselves as they removed their personal protective equipment (PPE). Many health care workers apparently lacked a clear understanding of how best to remove PPE without contaminating themselves. However, the correct order of removal as presented in the report was extremely condensed (gloves first, followed by mask and goggles).¹

In fact, little information about the appropriate sequence of removing PPE is available, and what is available often contains contrasting recommendations (Table 1).²⁻⁶ Moreover, for several of these sets of recommendations, hands potentially contaminated through con-

tact with patients' droplets and secretions present on the PPE could contact the nose, mouth or eyes while the health care worker is removing his or her mask and eye protection.

Careful hand hygiene plays a pivotal role in reducing the risk of transmission of SARS. Accordingly, the National Institute for Infectious Diseases in Italy has developed a procedure whereby the health care worker removes most PPE while wearing mask and eye protection and carefully decontaminates the hands before removing protection of the mucous membranes of the face. This procedure must be carried out in the health care worker's change area, outside the patient's isolation room.

Vincenzo Puro
Emanuele Nicastrì

National Institute for Infectious Diseases
IRCCS Lazzaro Spallanzani
Rome, Italy

References

1. Cluster of severe acute respiratory syndrome cases among protected health care workers — Toronto, April 2003. *Can Commun Dis Rep* 2003;11:93-7.
2. Young JG, D' Cunha C. Directive 03-05(R). *Directives to all Ontario acute care hospitals regarding infection control measures for SARS units*. Toronto: Ministry of Health and Long-Term Care; 2003 Apr 24. Available: www.homecareontario.ca/docs/sars/Apr%2024%20-%20Infection%20Control%20Directive%20Acute%20Care%20Hospitals.pdf (accessed 2003 Nov 6).
3. *Interim guidelines for national SARS preparedness*. Manila, Philippines: World Health Organization, Western Pacific Regional Office; updated

2003 May 13, revised 2003 May 26. Available: www.wpro.who.int/sars/interim_guidelines/interim_guidelines_26May.pdf (accessed 2003 Nov 6).

4. *Interim Australian infection control guidelines for severe acute respiratory syndrome (SARS)*. Canberra: Australian Government Department of Health and Ageing; updated 2003 May 16. Available: www.health.gov.au/sars/guidelines/index.htm (accessed 2003 Nov 6).
5. Ahuja AT, Wong JKT. Management and infection control in a radiology department during the SARS outbreak. In: *Radiological appearance of recent cases of atypical pneumonia in Hong Kong*. Shatin, NT, Hong Kong: Prince of Wales Hospital; 2003 Mar 21, updated 2003 Jul 30. Available: www.droid.cuhk.edu.hk/web/atypical_pneumonia/atypical_pneumonia.htm#infection (accessed 2003 Nov 6). See "Specific personal infection control guidelines for staff and patients."
6. Ng PC, So KW, Leung TF, Cheng FW, Lyon DJ, Wong W, et al. Infection control for SARS in a tertiary neonatal centre. *Arch Dis Child Fetal Neonatal Ed* 2003;88(5):F405-9.

DOI:10.1053/cmaj.1031700

Rehabilitation in developing countries

As Mohan Radhakrishna and Lalith Satkunam point out in their introduction to the recent *CMAJ* series on rehabilitation,¹ this important topic is not covered appropriately in medical schools. Furthermore, rehabilitation of those affected by physical and mental disability is neglected in developing countries, often as a result of corruption, poor management of financial resources and lack of natural resources.

Almost 10% of the global population is disabled in some way,² and most

Table 1: Procedures for removing personal protective equipment worn while caring for patients with SARS

Order	Ontario Ministry of Health and Long-Term Care ²	WHO, Western Pacific Regional Office ³	Australian Government Department of Health and Ageing ⁴	Department of Diagnostic Imaging, Prince of Wales Hospital, Hong Kong ⁵	Tertiary neonatal centre, Prince of Wales Hospital, Hong Kong ⁶	National Institute for Infectious Diseases, Rome, Italy
1	Gloves*	Wash gloved hands	Gloves	Gloves	Hair and shoe covers, gown	Hair and shoe covers, gown
2	Hand hygiene†	Aprons, gowns and shoe covers	Eye protection	Hair cover	Gloves	Gloves
3	Eye protection	Gloves	Gown	Eye protection	Hand hygiene	Hand hygiene
4	Mask	Goggles	Mask	Gown	Eye protection	Eye protection
5	Hair cover	Hair cover and mask	Hand hygiene	Mask	Mask	Hand hygiene
6	Hand hygiene	Hand hygiene		Hand hygiene	Hand hygiene	Mask
7	—	—	—	—	—	Hand hygiene

Note: WHO = World Health Organization.

*Second pair. First pair and gown to be removed soon after providing direct patient care, while still inside the patient's room.

†Hand-washing or hand-rubbing, or both, with alcohol or other antiseptic agents.