

This article has been peer reviewed.

Jonathan Klein is Director of the American Academy of Pediatrics Julius B. Richmond Center of Excellence for Children and Secondhand Smoke and is Associate Professor of Pediatrics and of Community and Preventive Medicine, University of Rochester School of Medicine, Rochester, NY.

Competing interests: None declared.

Acknowledgement: Jonathan Klein's work on this commentary was supported in part by the Flight Attendant Medical Research Institute and by the American Legacy Foundation.

REFERENCES

1. World Health Organization. Statement from Catherine Le Galès — Camus, Assistant Director-General, Non Communicable Diseases and Mental Health. Available: www.who.int/tobacco/communications/events/wntd/2006/statement/en/index.html (accessed 2006 Jun 27).
2. US Centers for Disease Control and Prevention. Trends in the prevalence of cigarette use. Available: www.cdc.gov/HealthyYouth/yrbs/pdf/trends/2005_YRBS

3. Gervais A, O'Loughlin J, Meshfedjian G, et al. Milestones in the natural course of onset of cigarette use among adolescents. *CMAJ* 2006;175(3):255-61.
4. DiFranza JR, Wellman RJ, Sargent JD, et al; Tobacco Consortium, Center for Child Health Research of the American Academy of Pediatrics. Tobacco promotion and the initiation of tobacco use: assessing the evidence for causality. *Pediatrics* 2006; 117:e1237-48.
5. DiFranza JR, Savageau JA, Fletcher K, et al. Measuring the loss of autonomy over nicotine use in adolescents: the DANDY (Development and Assessment of Nicotine Dependence in Youths) study. *Arch Pediatr Adolesc Med* 2002;156:397-403.
6. Fiore MC, Bailey WC, Cohen SJ, et al. *Treating tobacco use and dependence*. Clinical Practice Guideline. Rockville (MD): Public Health Service, US Department of Health and Human Services; June 2000. Available: www.surgeongeneral.gov/tobacco/treating_tobacco_use.pdf (accessed 2006 June 29).
7. Pbert L, Moolchan ET, Muramoto M, et al; Tobacco Consortium, Center for Child Health Research of the American Academy of Pediatrics. The state of office-based interventions for youth tobacco use. *Pediatrics* 2003;111:e650-60.

Correspondence to: Dr. Jonathan D. Klein, University of Rochester School of Medicine, 690–601 Elmwood Ave., Rochester NY 14642, USA; fax 585 242-9733; jonathan_klein@urmc.rochester.edu

Personal protective equipment for preventing respiratory infections: What have we really learned?

John M. Conly

∞ See related article page 249

The use of personal protective equipment (PPE) for health care workers (HCWs) has evolved from the isolation precautions first implemented years ago for patients with communicable diseases such as smallpox, tuberculosis and diphtheria.¹ The use of PPE (gloves, gowns, masks and eye protection) in combination with single rooms with airflow control represents the usual barrier precautions employed to prevent transmission of pathogenic microorganisms to HCWs. The mechanisms of transmission (airborne, droplet, contact, vector or common vehicle) for the microbe in question often mandate the specific combination of barrier precautions chosen.²

Reports of SARS among HCWs in hospital outbreaks reported from Canada, China, Hong Kong, Taiwan and Vietnam focused attention on the critical importance of infection-control practices, including the use of PPE, and the role of training and knowledge among HCWs in using PPE and barrier precautions appropriately.³ Microbes transmitted by the airborne or droplet routes create the greatest anxiety among HCWs. Additional risks for transmission are posed by the emergence of new pathogens with a severe illness profile (e.g., SARS and avian influenza) and immuno- and other highly compromised patients, who may carry greater microbial burdens for prolonged periods. The advent of new technological diagnostic and therapeutic modalities may also lengthen HCWs' exposure to patients carrying highly infective pathogens.

A thorough understanding of the usual routes of transmission of microbes and the conditions under which these routes may change is paramount to prevent the spread of an infection.² Contact transmission, the most common route, occurs when microbes are transferred either directly by physical contact between an infected or colonized individual and a new host or indirectly via an intermediate object (a fomite).² Droplet transmission involves drops of fluid 5 µm

SARS focused attention on the critical importance of infection-control practices.

in diameter and larger, produced from the respiratory tract during coughing or sneezing or by medical procedures, propelled within 1 m of the source patient. Airborne transmission refers to dissemination of microbes within droplet nuclei (particles < 5 µm in diameter), which result from the evaporation of larger droplets or exist within dust particles and remain suspended in the air for long periods. Although most respiratory viruses are transmitted by droplet and contact methods, microbes that can spread via airborne trans-

mission include the agents of measles, smallpox, tuberculosis and varicella-zoster.

The SARS outbreaks helped us to recognize the enhanced transmissibility of respiratory pathogens during respiratory procedures that may generate aerosol particles. These procedures have the potential to generate a multitude of large and small droplets, and the procedure itself may propel these droplets well beyond the 1-m radius usually associated with larger droplets. Agreement about aerosol-generating procedures is not universal, but the use of nebulizers, high-flow oxygen, bronchoscopy, non-intubated ventilation (continuous or bilevel positive airway pressure), bag-valve ventilation and uncontrolled intubation are considered higher-risk procedures;⁴ they can cause the lines between droplet and airborne transmission to become blurred. What SARS has taught us is that the use of these specialized respiratory procedures can increase the potential for episodic localized airborne transmission and probably expand opportunities for fomite and droplet transmission.

There is compelling evidence that the SARS coronavirus is spread through droplet and contact transmission.³ Early reports of high infection rates among HCWs and so-called superspreading events were incorrectly judged to indicate a high level of communicability and led to an assumption that the pathogen was airborne.^{5,6} Patients with unrecognized SARS, inadequate understanding among HCWs, a lack of compliance with basic infection-control measures and the creation of virus-laden aerosols provide the best explanation for the nosocomial outbreaks of SARS.^{5,7,8} Although some HCWs were reported to have become infected with SARS despite the use of PPE, most of these infections occurred during high-risk aerosol- and droplet-generating procedures, accompanied by accounts of suboptimal compliance with protocols for the donning or removal of PPE, PPE reuse, inappropriate double-gloving and gowning (with potential cross-contamination), fatigue and poor knowledge of basic procedures for infection control, which may provide explanations for transmission.^{3,8,9}

The report by Zamora and colleagues¹⁰ in this issue of *CMAJ* illustrates the potential for contamination (which represents a potential for contact transmission) with the use of 2 different personal protective systems: a standard procedure with gloves, gowns, masks and eye protection, or one that incorporates a more elaborate powered air-purifying respirator (PAPR). They conducted a well-designed crossover analysis with adequate power to detect significant outcome differences in base-clothing or skin contamination, using a standard protocol in a controlled setting and a suitable surrogate marker for contamination. They found that skin contamination with the surrogate marker occurred with either PPE system; exposed skin contamination occurred more often with standard PPE than with the PAPR system; and PPE donning and removal violations occurred more often with use of the PAPR system.

Both systems have their faults and may create potential risks for contact transmission, either through direct contamination or when donning and removal protocols are breached.

Although the study begs the question as to how applicable these results would be in an uncontrolled real-life scenario, it certainly emphasizes the need for handwashing after glove removal, given the high contamination rates of the hands and wrists with the use of either system. They also provide indirect evidence that whatever system is used, the need for trials, drills and adherence to protocol are important elements in the protection of HCWs. Any system or strategy can be expected to meet with success, but execution becomes a critical element in the overall process. The consistent application of appropriate infection-control techniques is essential to the prevention of droplet and contact transmission. This has been demonstrated in many countries around the world,³ most of which had no access to PAPR systems and many even to N95 respirator masks, but were nevertheless able to focus on adherence to infection-control techniques, which was the key component in controlling the spread of SARS.

This article has been peer reviewed.

John Conly is with the Divisions of Infectious Diseases and Microbiology, Departments of Medicine, Pathology and Laboratory Medicine and of Microbiology and Infectious Diseases, University of Calgary and the Calgary Health Region, Calgary, Alta.

Competing interests: None declared.

Acknowledgements: I thank Dr. Manuel Mah and Karen Hope for their helpful comments and critique of this commentary.

REFERENCES

1. Jackson M, Lynch P. Isolation practices: a historical perspective. *Am J Infect Contr* 1985;13:21-31.
2. Health Canada, Laboratory Centre for Disease Control, Bureau of Infectious Diseases, Division of Nosocomial and Occupational Infections. Routine practices and additional precautions for preventing the transmission of infection in health care. *Can Commun Dis Rep* 1999; 25(Suppl 4): S1-142. Available: www.phac-aspc.gc.ca/publicat/ccdr-rmtc/g99pdf/cdr25s4e.pdf (accessed 2006 June 12).
3. Shaw K. The 2003 SARS outbreak and its impact on infection control practices. *Public Health* 2006;120:8-14.
4. Health Canada. Infection control precautions for respiratory infections transmitted by large droplet and contact: infection control guidance if there is a SARS outbreak anywhere in the world, when an individual presents to a health care institution with a respiratory infection [draft]. 2003 Dec 17. Available: www.phac-aspc.gc.ca/sars-sras/pdf/sars-icg-outbreakworld_e.pdf (accessed 2006 Jun 12).
5. Varia M, Wilson S, Sarwal S, et al. Investigation of a nosocomial outbreak of severe acute respiratory syndrome (SARS) in Toronto, Canada. *CMAJ* 2003;169(4):285-92.
6. Shen Z, Ning F, Zhou W, et al. Superspreading SARS events, Beijing, 2003. *Emerg Infect Dis* 2004;10:256-60.
7. Lau JT, Fung KS, Wong TW, et al. SARS transmission among hospital workers in Hong Kong. *Emerg Infect Dis* 2004;10:280-6.
8. Gamage D, Moore R, Copes R, et al; The BC Interdisciplinary Respiratory Protection Study Group. Protecting health care workers from SARS and other respiratory pathogens: a review of the infection control literature. *Am J Infect Control* 2005; 33:114-21.
9. Ofner-Agnostini M, Gravel D, McDonald C, et al. Cluster of cases of severe acute respiratory syndrome among Toronto healthcare workers after implementation of infection control precautions: a case series. *Infect Control Hosp Epidemiol* 2006; 27:473-8.
10. Zamora JE, Murdoch J, Simchison B, et al. Contamination: a comparison of 2 personal protective systems. *CMAJ* 2006;175(3):249-54.

Correspondence to: Dr. John Maynard Conly, University of Calgary, North Tower, Rm. 930, Foothills Medical Centre, 1403 - 29th St. NW, Calgary AB T2N 2T9; fax 403 922-1095; john.conly@calgaryhealthregion.ca