

**Appendix 2 (as submitted by author). Input parameter estimates and standard deviations for model**

Variable	Input parameter estimate	Standard deviation	Source
Prevalence of low risk polyp at 50 years	0.089	0.009	W1-W5
Prevalence of advanced adenoma at 50 years	0.03	0.003	W1-W5
Prevalence of pre-clinical cancer at 50 years	0.003	0.0003	W1-W5
<b>Annual transition probabilities</b>			
Low risk polyp from normal	Age-dependent	± 10% input parameter	W1-W5
Advanced adenoma from low risk polyp	0.036	0.011	W1-W5
Localized cancer from advanced adenoma	0.042	0.013	W1-W5
Regional cancer from localized cancer	0.17	0.052	W1,W5-W7
Distant cancer from regional cancer	0.10	0.029	W1,W3,W5,W7, W8
Diagnosed from pre-clinical localized cancer	0.17	0.050	W1,W5-W7
Diagnosed from pre-clinical regional cancer	0.21	0.064	W1,W5-W7
Diagnosed from pre-clinical distant cancer	1	--	Assumption
<b>Screening test performance</b>			
<b>Low sensitivity guaiac fecal occult blood test</b>			
Sensitivity to detect a low risk polyp	0.05	0.02	W8-W11
Sensitivity to detect an advanced adenoma	0.19	0.05	W8-W11
Sensitivity to detect cancer	0.46	0.31	W8-W11
Specificity	0.94	0.05	W8-W11
<b>High sensitivity guaiac fecal occult blood test</b>			
Sensitivity to detect a low risk polyp	0.03	0.04	W11,W12
Sensitivity to detect an advanced adenoma	0.34	0.17	W11-W13
Sensitivity to detect cancer	0.72	0.32	W11-W13
Specificity	0.91	0.17	W11-W13
<b>Fecal immunochemical test</b>			
Sensitivity to detect a low risk polyp	0.10	0.11	W11,W12,W14
Sensitivity to detect an advanced adenoma	0.46	0.13	W11-W16
Sensitivity to detect cancer	0.85	0.10	W11-W16
Specificity	0.94	0.26	W11-W15
<b>Fecal DNA</b>			
Sensitivity to detect a low risk polyp	0.08	0.008	W10
Sensitivity to detect an advanced adenoma	0.45	0.28	W17-W19
Sensitivity to detect cancer	0.70	0.12	W17-W19
Specificity	0.94	0.06	W10,W18,W19
<b>Double contrast barium enema</b>			
Sensitivity to detect a low risk polyp	0.35	0.07	W20,W21
Sensitivity to detect an advanced adenoma	0.47	0.03	W20,W21
Sensitivity to detect cancer	0.86	0.04	W20,W22
Specificity	0.82	0.082	W20
<b>Colonoscopy and sigmoidoscopy (distal only)</b>			
Probability of a negative sigmoidoscopy and a proximal neoplasm	0.21	0.10	W2
Sensitivity to detect a low risk polyp	0.92	0.09	W22-W28
Sensitivity to detect an advanced adenoma	0.97	0.05	W22-W29
Sensitivity to detect cancer	0.93	0.19	W22-W29
Specificity	1	-	W20, W23,W27

<b>CT colonography</b>			
Sensitivity to detect a low risk polyp	0.50	0.25	W20, W23,W24,W26, W28,W30,W31
Sensitivity to detect an advanced adenoma	0.78	0.19	W20,W23,W24, W26,W30,W31
Sensitivity to detect cancer	0.89	0.12	W20,W23,W24,W 26,W28,W31
Specificity	0.91	0.07	W20,W23,W24, W26,W28,W31
<b>Compliance</b>			
Probability of undergoing each screening round	0.73	0.22	W32-W35
Follow-up colonoscopy for a positive screening test	0.84	0.25	W35-W37
<b>Serious complication</b>			
Sigmoidoscopy	0.0002	0.00006	W38
Colonoscopy diagnostic	0.0008	0.0002	W39
Colonoscopy with polypectomy	0.007	0.0021	W39
Double contrast barium enema	0.0002	0.00006	W40
CT colonography	0.0001	0.00003	W41
Death from complication	0.012	0.0036	W39
<b>Relapse (annual)</b>			
Stage I	0.01	0.003	W42
Stage II/III	0.04	0.012	W43
<b>Medical costs (2007 Canadian dollars)</b>			
Primary Care Physician Visit	58	17	W44
Guaiac-based fecal occult blood test	10	3	W45
Fecal immunochemical test	25	7	W46
Fecal DNA	346	92	W46
Sigmoidoscopy	193	58	W47,W48
Double contrast barium enema	178	53	W47
Colonoscopy	424	128	W47,W48
Polypectomy	185	56	W47
CT colonography	590	177	W47
Complication	31,193	9,358	W49
<b>Cost of colorectal cancer (2007 Canadian dollars)</b>			
Stage I			
Year 1	13,734	4,120	W50
Year 2-5 (annual)	503	151	W51
Stage II			
Year 1	27,095	8,129	W50
Year 2-5 (annual)	857	257	W51
Stage III			
Year 1	35,040	10,512	W50
Year 2-5 (annual)	857	257	W51
Stage IV			
Year 1	72,281	21,684	W50
Year 2-5 (annual)	670	201	W51
Relapse (annual)	11,435	3,431	W50
Terminal Care	17,869	5,361	W50
<b>Utilities</b>			
No colorectal cancer	1.0	0	Assumption
Stage I colorectal cancer	0.74	0.22	W52
Stage II colorectal cancer	0.69	0.21	W52
Stage III colorectal cancer	0.64	0.19	W52
Stage IV colorectal cancer	0.25	0.075	W52
Colorectal cancer relapse	0.69	0.21	Assumption
Death	0.00	0	Assumption

### References for model

- W1. Regula J, Rupinski M, Kraszewska E, et al. Colonoscopy in colorectal-cancer screening for detection of advanced neoplasia. *N Engl J Med* 2006;355:1863-72.
- W2. Strul H, Kariv R, Leshno M, et al. The prevalence rate and anatomic location of colorectal adenoma and cancer detected by colonoscopy in average-risk individuals aged 40-80 years. *Am J Gastroenterol* 2006;101:255-62.
- W3. Schoenfeld P, Cash B, Flood A, et al. Colonoscopic screening of average-risk women for colorectal neoplasia. *N Engl J Med* 2005;352:2061-8.
- W4. Lin OS, Kozarek RA, Schembre DB, et al. Screening colonoscopy in very elderly patients: prevalence of neoplasia and estimated impact on life expectancy. *JAMA* 2006;295:2357-65.
- W5. Imperiale TF, Wagner DR, Lin CY, et al. Results of screening colonoscopy among persons 40 to 49 years of age. *N Engl J Med* 2002;346:1781-5.
- W6. Lieberman DA, Weiss DG, Bond JH, et al. Use of colonoscopy to screen asymptomatic adults for colorectal cancer. Veterans Affairs Cooperative Study Group 380. *N Engl J Med* 2000;343:162-8.
- W7. Surveillance, epidemiology, and end results (SEER) program. SEER\*Stat Database: Incidence — SEER 9 Regs Public Use, Nov. 2003 Sub (1973–2004). National Cancer Institute, DCCPS, Surveillance Research Program; 2004. Available: [www.seer.cancer.gov](http://www.seer.cancer.gov) (accessed 2008 Aug. 12).
- W8. Collins JF, Lieberman DA, Durbin TE, et al. Accuracy of screening for fecal occult blood on a single stool sample obtained by digital rectal examination: a comparison with recommended sampling practice. *Ann Intern Med* 2005;142:81-5.
- W9. Lieberman DA, Weiss DG. One-time screening for colorectal cancer with combined fecal occult-blood testing and examination of the distal colon. *N Engl J Med* 2001;345:555-60.
- W10. Imperiale TF, Ransohoff DF, Itzkowitz SH, et al. Fecal DNA versus fecal occult blood for colorectal-cancer screening in an average-risk population. *N Engl J Med* 2004;351:2704-14.
- W11. Greenberg PD, Bertario L, Gnauck R, et al. A prospective multicenter evaluation of new fecal occult blood tests in patients undergoing colonoscopy. *Am J Gastroenterol* 2000;95:1331-8.
- W12. Hoepffner N, Shastri YM, Hanisch E, et al. Comparative evaluation of a new bedside faecal occult blood test in a prospective multicentre study. *Aliment Pharmacol Ther* 2006;23:145-54.
- W13. Wong BC, Wong WM, Cheung KL, et al. A sensitive guaiac faecal occult blood test is less useful than an immunochemical test for colorectal cancer screening in a Chinese population. *Aliment Pharmacol Ther* 2003;18:941-6.
- W14. Shastri YM, Loitsch S, Hoepffner N, et al. Comparison of an established simple office-based immunological FOBT with fecal tumor pyruvate kinase type M2 (M2-PK) for colorectal cancer screening: prospective multicenter study. *Am J Gastroenterol* 2008;103:1496-504.
- W15. Levi Z, Rozen P, Hazazi R, et al. A quantitative immunochemical fecal occult blood test for colorectal neoplasia. *Ann Intern Med* 2007;146:244-55.
- W16. Yoshinaga M, Motomura S, Takeda H, et al. Evaluation of the sensitivity of an immunochemical fecal occult blood test for colorectal neoplasia. *Am J Gastroenterol* 1995;90:1076-9.
- W17. Syngal S, Stoffel E, Chung D, et al. Detection of stool DNA mutations before and after treatment of colorectal neoplasia. *Cancer* 2006;106:277-83.
- W18. Ahlquist DA, Skoletsky JE, Boynton KA, et al. Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a multitarget assay panel. *Gastroenterology* 2000;119:1219-27.
- W19. Tagore KS, Lawson MJ, Yucaitis JA, et al. Sensitivity and specificity of a stool DNA multitarget assay panel for the detection of advanced colorectal neoplasia. *Clin Colorectal Cancer* 2003;3:47-53.
- W20. Rockey DC, Paulson E, Niedzwiecki D, et al. Analysis of air contrast barium enema, computed tomographic colonography, and colonoscopy: prospective comparison. *Lancet* 2005;365:305-11.
- W21. Winawer SJ, Stewart ET, Zauber AG, et al. A comparison of colonoscopy and double-contrast barium enema for surveillance after polypectomy. National Polyp Study Work Group. *N Engl J Med* 2000;342:1766-72.
- W22. Irvine EJ, O'Connor J, Frost RA, et al. Prospective comparison of double contrast barium enema plus flexible sigmoidoscopy v colonoscopy in rectal bleeding: barium enema v. colonoscopy in rectal bleeding. *Gut* 1988;29:1188-93.
- W23. Cotton PB, Durkalski VL, Pineau BC, et al. Computed tomographic colonography (virtual colonoscopy): a multicenter comparison with standard colonoscopy for detection of colorectal neoplasia. *JAMA* 2004;291:1713-9.

- W24. Pickhardt PJ, Choi JR, Hwang I, et al. Computed tomographic virtual colonoscopy to screen for colorectal neoplasia in asymptomatic adults. *N Engl J Med* 2003;349:2191-200.
- W25. Rex DK, Cutler CS, Lemmel GT, et al. Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies. *Gastroenterology* 1997;112:24-8.
- W26. Ginnerup Pedersen B, Christiansen TE, Bjerregaard NC, et al. Colonoscopy and multidetector-array computed-tomographic colonography: detection rates and feasibility. *Endoscopy* 2003;35:736-42.
- W27. Hixson LJ, Fennerty MB, Sampliner RE, et al. Prospective blinded trial of the colonoscopic miss-rate of large colorectal polyps. *Gastrointest Endosc* 1991;37:125-7.
- W28. Taylor SA, Halligan S, Saunders BP, et al. Use of multidetector-row CT colonography for detection of colorectal neoplasia in patients referred via the Department of Health "2-Week-wait" initiative. *Clin Radiol* 2003;58:855-61.
- W29. Matsushita M, Hajiro K, Okazaki K, et al. Efficacy of total colonoscopy with a transparent cap in comparison with colonoscopy without the cap. *Endoscopy* 1998;30:444-7.
- W30. Rex DK, Alikhan M, Cummings O, et al. Accuracy of pathologic interpretation of colorectal polyps by general pathologists in community practice. *Gastrointest Endosc* 1999;50:468-74.
- W31. Pineau BC, Paskett ED, Chen GJ, et al. Virtual colonoscopy using oral contrast compared with colonoscopy for the detection of patients with colorectal polyps. *Gastroenterology* 2003;125:304-10.
- W32. Hardcastle JD, Chamberlain JO, Robinson MH, et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996;348:1472-7.
- W33. Kronborg O, Fenger C, Olsen J, et al. Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet* 1996;348:1467-71.
- W34. Mandel JS, Bond JH, Church TR, et al. Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N Engl J Med* 1993;328:1365-71.
- W35. Selby JV, Friedman GD, Quesenberry CP Jr, et al. A case-control study of screening sigmoidoscopy and mortality from colorectal cancer. *N Engl J Med* 1992;326:653-7.
- W36. Jorgensen OD, Kronborg O, Fenger C. A randomised study of screening for colorectal cancer using faecal occult blood testing: results after 13 years and seven biennial screening rounds. *Gut* 2002;50:29-32.
- W37. Mandel JS, Church TR, Bond JH, et al. The effect of fecal occult-blood screening on the incidence of colorectal cancer. *N Engl J Med* 2000;343:1603-7.
- W38. Levin TR, Conell C, Shapiro JA, et al. Complications of screening flexible sigmoidoscopy. *Gastroenterology* 2002;123:1786-92.
- W39. Levin TR, Zhao W, Conell C, et al. Complications of colonoscopy in an integrated health care delivery system. *Ann Intern Med* 2006;145:880-6.
- W40. de Feiter PW, Soeters PB, Dejong CH. Rectal perforations after barium enema: a review. *Dis Colon Rectum* 2006;49:261-71.
- W41. Sosna J, Blachar A, Amitai M, et al. Colonic perforation at CT colonography: assessment of risk in a multicenter large cohort. *Radiology* 2006;239:457-63.
- W42. Weiser MR, Landmann RG, Kattan MW, et al. Individualized prediction of colon cancer recurrence using a nomogram. *J Clin Oncol* 2008;26:380-5.
- W43. de Gramont A, Boni C, Navarro M, et al. Oxaliplatin/5FU/LV in adjuvant colon cancer: updated efficacy results of the MOSAIC trial including survival, with a median follow-up of six years. *J Clin Oncol* 2007;25(18S):AB4007.
- W44. Ontario health insurance schedule of benefits and fees. Toronto (ON): Ontario Ministry of Health; 2007.
- W45. Flanagan WM, Le Petit C, Berthelot JM, et al. Potential impact of population-based colorectal cancer screening in Canada. *Chronic Dis Can* 2003;24:81-8.
- W46. Parekh M, Fendrick AM, Ladabaum U. As tests evolve and costs of cancer care rise: reappraising stool-based screening for colorectal neoplasia. *Aliment Pharmacol Ther* 2008;27:697-712.
- W47. McGrath JS, Ponich TP, Gregor JC. Screening for colorectal cancer: the cost to find an advanced adenoma. *Am J Gastroenterol* 2002;97:2902-7.
- W48. Sambrook JC, Chui W, Wang H, et al. Colonoscopy and flexible sigmoidoscopy: A microcosting study evaluating diagnostic procedural costs in an outpatient endoscopy clinic. *Can J Gastroenterol* 2006;20(Suppl A):AB167.

- W49. Heitman SJ, Manns BJ, Hilsden RJ, et al. Cost-effectiveness of computerized tomographic colonography versus colonoscopy for colorectal cancer screening. *CMAJ* 2005;173:877-81.
- W50. Maroun J, Ng E, Berthelot JM, et al. Lifetime costs of colon and rectal cancer management in Canada. *Chronic Dis Can* 2003;24:91-101.
- W51. Figueiredo A, Rumble RB, Maroun J, et al. Follow-up of patients with curatively resected colorectal cancer: a practice guideline. *BMC Cancer* 2003;3:26.
- W52. Ness RM, Holmes AM, Klein R, et al. Utility valuations for outcome states of colorectal cancer. *Am J Gastroenterol* 1999;94:1650-7.