

Appendix 6 (as supplied by the authors): Subgroup analysis of Risk of Myocardial Infarction Among Patients Using ARBs and ACEIs

Characteristic	Hazard Ratio [*] (95% CI)	P Value	Interaction P Value
Gender			
Male	1.009 (0.831-1.225)	0.9	0.8
Female	1.052 (0.805-1.375)	0.7	
Age			
20-64 years	1.105 (0.907-1.348)	0.3	0.2
≥65 years	0.903 (0.698-1.170)	0.4	
Charlson Comorbidity Index score			
2	1.323 (0.974-1.798)	0.07	0.1
3	1.049 (0.752-1.463)	0.8	
4	1.067 (0.726-1.568)	0.7	
≥5	0.818 (0.625-1.072)	0.1	
Hypertension			
Yes	1.033 (0.877-1.217)	0.7	0.8
No	0.963 (0.556-1.668)	0.9	
Chronic kidney disease			
Yes	0.911 (0.650-1.277)	0.59	0.4
No	1.062 (0.889-1.268)	0.5	
Heart failure			
Yes	0.856 (0.542-1.351)	0.5	0.4
No	1.051 (0.889-1.242)	0.6	
Coronary artery disease			
Yes	0.914 (0.692-1.208)	0.5	0.3
No	1.085 (0.897-1.313)	0.4	

* Adjusted for propensity score; ACEI cohort: reference group

Abbreviations: CI, confidence interval; ARBs, Angiotensin II receptor blockers; ACEIs, angiotensin-converting-enzyme inhibitors.